

## New England Biolabs Certificate of Analysis

**Product Name:** *rCutSmart™ Buffer*  
**Catalog Number:** *B6004S*  
**Concentration:** *10 X Concentrate*  
**Packaging Lot Number:** *10208664*  
**Expiration Date:** *09/2026*  
**Storage Temperature:** *-20°C*  
**Specification Version:** *PS-B6004S/V v1.0*  
**Composition (1X):** *50 mM Potassium Acetate, 20 mM Tris Acetate, 10 mM Magnesium Acetate, 100 µg/ml rAlbumin, (pH 7.9 @ 25°C)*

rCutSmart™ Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B6004SVIAL	rCutSmart™ Buffer	10204839	Pass

Assay Name/Specification	Lot # 10208664
<b>Conductivity (buffers/solutions)</b> The conductivity of 10X rCutSmart™ Buffer is between 40 and 46 mS/cm at 25°C.	Pass
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Functional Testing (Restriction Digest, Buffer)</b> A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and 1 unit of ClaI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
<b>Functional Testing (Restriction Digest, Buffer)</b> A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of EcoRI-HF® incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
<b>Functional Testing (Restriction Digest, Buffer)</b> A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass

Assay Name/Specification	Lot # 10208664
<p><b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>RNase Activity (Buffer)</b> A 10 µl reaction in 1X rCutSmart™ Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>
<p><b>pH (buffers/solutions)</b> The pH of 10X rCutSmart™ Buffer is between pH 7.8 and 8.0 at 25°C.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Nancy Considine  
Production Scientist  
26 Sep 2023



Josh Hersey  
Packaging Quality Control Inspector  
02 Oct 2023