

New England Biolabs Certificate of Analysis

Product Name: BsmBI-v2
Catalog Number: R0739L
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 55°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10059568
Expiration Date: 11/2021
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)
Specification Version: PS-R0739S/L v1.0

BsmBI-v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0739LVIAL	BsmBI-v2	10058638	Pass
B7203SVIAL	NEBuffer™ 3.1	10053972	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10046085	Pass

Assay Name/Specification	Lot # 10059568
Functional Testing (15 minute Digest) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and 1 µl of BsmBI-v2 incubated for 15 minutes at 55°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BsmBI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsmBI-v2.	Pass
Protein Purity Assay (SDS-PAGE) BsmBI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of BsmBI-v2 incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.	Pass

Assay Name/Specification	Lot # 10059568
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of BsmBI-v2 incubated for 4 hours at 55°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 10 units of BsmBI-v2 incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Jianying Luo
Production Scientist
01 Nov 2019



Michael Tonello
Packaging Quality Control Inspector
13 Nov 2019